



Lab 5: Properties of a Group in the Periodic Table



Goals

1. To introduce the concept of element groups
2. To show how elements in their groups are similar to one another and to explain why this is so

Materials and Equipment

Slide of elements:

Carbon
Silicon
Germanium
Tin

Digital voltmeter

Materials Not Included

Periodic table
Ice
Hot water
Shallow dish

Introduction

The periodic table is periodic, which means that it repeats. This happens because atoms fill their outer-shell electron orbits in an organized way. As it turns out, the outer-shell electrons are almost entirely responsible for the chemical properties of the elements.

The columns on the periodic table contain elements that have identical outer-shell electrons. These columns are called element groups. Group I elements are comprised of the alkali metals, and they are all very reactive. They form ionic compounds with many elements that are in

the group VI and group VII columns. Other groups have other similarities.

In this laboratory we will study the group IVB elements, which are carbon, silicon, germanium, tin, and lead. We will omit lead from the actual hands-on examination. All the group IVB elements are quite commonly seen in ordinary life. Elemental carbon can be seen if you badly burn your toast or in the pencil "lead" you write with. Silicon is the basis for most electronic microchips in use today. Germanium is less common, but it too is used to make microchips. Tin is used in most solders. Lead is a solder, as well as a weight for fishing and balancing car tires.

Many of the group IVB elements are classified as semiconductors. They are not insulators since they can pass some current, but they are not conductors since they have a higher resistance than most metals. Semiconductors have an electrical resistance that is sensitive to temperature. You will investigate their changing resistance in this laboratory.

Devotional

"When I consider your heavens, the work of your fingers, the moon and the stars, which you have set in place, what is man that you are mindful of him, the son of man that you care for him? You made him a little lower than the heavenly beings and crowned him with glory and honor. You made him ruler over the works of your hands; you put everything under his feet ... O LORD, our Lord, how majestic is your name in all the earth! Ps. 8:3-6, 9

Principle: God's world is surprisingly well organized. When we recognize this, it may bring a sense of wonder.

From the heavens above, to the seasons, to the migrating birds, to the smallest particles, all creation shows forth God's organization. The periodic table is clear evidence of God's organized patterns in the world of chemistry. The organization of the table is due to electrons filling their outer-shell orbitals. The groups, which are found in columns in the periodic table, all have similar properties. The elements on the left side weakly hold their electrons, while those on the right side fight to gain more electrons.

The metals all have similar "metallic" properties. The rare earth elements, such as the lanthanide series metals, fill their f-shell, which makes them so similar to each other that they are almost impossible to separate. The patterns are clear. And for all of us who see God's hand in these patterns, the elegance of the periodic table can bring a sense of wonder. How wonderful are your works our LORD!

Procedure

1. Using a periodic table, write out the electronic structures of all the group IVB elements. Put your answers into Table 1.
2. Check the room temperature resistance of *C*, *Si*, *Ge*, and *Sn*. To do this you will put the multimeter on the resistance range in the 200 Ohm scale. Touch one probe to each end of each sample provided. Wait until the value is relatively stable and report an approximate value. The values do move around a bit, so make your best estimate. Report the values in Table 2.
3. Make some ice water and pour it into a very shallow dish. Place the slide with the sample elements into the ice water and wait a minute for them to cool. Dry the samples and quickly measure the resistance of each sample. Again, you will need to wait for the resistance value to stabilize and then make your best estimate of the value. Record your results in Table 2.
4. Repeat the test again, using hot water in the shallow dish instead of cold water. You will need to preheat the dish with hot water to make sure the water remains as hot as possible.
5. Take a knife or small screwdriver and attempt to make a very small scratch on the element samples. If you have access to a lead wheel weight or a lead fishing weight, you may examine that and add the results to the table. Try to determine if the samples tend to be hard and brittle, or soft and ductile. Fill out column 3 in Table 1.

Lab 5

Questions for Properties of a Group in the Periodic Table



Table 1

Element	Electronic Structure	Brittle/Ductile/In Between
Carbon		
Silicon		
Germanium		
Tin		
Lead		

Lab
5

Table 2

Element	Resistance, In Ice Water	Resistance, Room Temperature	Resistance, In Hot Water
Carbon			
Silicon			
Germanium			
Tin			

1. In what ways are the electronic structures of the group IVB elements similar? In what ways are they different?

2. Compare the room temperature resistances for all the samples. Are they similar or quite different?

3. How does resistance change with temperature? Is there more resistance or less resistance at higher temperatures? Compare the change in resistance for all the samples.

4. Are the samples soft and ductile or hard and brittle? Are some in between?

5. What do you think causes the difference between the elements on the top of the periodic table, like carbon, with those on the bottom of the periodic table, like tin or lead?



Lab 29: Cross-Linking of a Polymer



Goals

1. To observe the cross-linking of a polymer
2. To better understand the process of solidification of a material by cross-linking
3. To observe a non-Newtonian fluid
4. To better understand a hydrogen bond

Materials and Equipment

150-mL beaker
30-mL beaker
Polyvinyl alcohol (*PVA*)
Sodium tetraborate
Wooden stir stick
Measuring spoon, 1.0 mL

Materials Not Included

Paper plate
Food coloring (optional)
Source of heat


Introduction

Many common materials solidify by forming cross-links between molecules. Examples include epoxy and ordinary silicon caulk. Cross-linking is the bonding together of adjacent polymer molecules, often by smaller molecules of another material. Cross-linking occurs in all directions and usually forms a three-dimensional network. Polymers can have thousands of monomers


in a single atom, and one polymer molecule can cross-link to many other molecules.

Most vinyl polymers are not water soluble. Polyvinyl alcohol (*PVA*) is soluble, however, because it has (OH^-) hydroxyl groups. This allows the *PVA* to form a colloidal suspension in water because the hydroxyl (OH^-) groups are ready acceptors for hydrogen bonding. (See the introduction for Lab 26: Organic Chemistry Models for more information about hydrogen bonding.) This allows *PVA* to interact with polar water molecules and makes *PVA* soluble. Because of the hydrogen bonding, *PVA* molecules will also interact slightly with each other. This increases the viscosity in the solution, depending on the concentration.

Borate ions, $\text{B}(\text{OH})_4^-$, are formed in stepwise reactions when sodium tetraborate dissolves in water. The first step is the formation of boric acid:



Boric acid then accepts a hydroxide from water:



Borate ions, $\text{B}(\text{OH})_4^-$, can form hydrogen bonds with all four of its hydroxyl groups between adjacent *PVA* polymer molecules. This is a hydrogen bond cross-link. Although these hydrogen bonds are relatively strong, they are also dynamic, so they are constantly forming and dissociating. This gives slime the ability to flow or ooze while still having some strength and elasticity short-term.

Slime is a good example of a non-

Newtonian fluid. A non-Newtonian fluid is a fluid in which the viscosity changes with the applied strain rate. It stretches when pulled slowly, but breaks when pulled quickly. As a result, non-Newtonian fluids may not have a well-defined viscosity.

Devotional

“As you come to him, the living Stone—rejected by men but chosen by God and precious to him—you also, like living stones, are being built into a spiritual house to be a holy priesthood, offering spiritual sacrifices acceptable to God through Jesus Christ.”

1 Peter 2:4-5

“Dear friends, let us love one another, for love comes from God. Everyone who loves has been born of God and knows God.” 1 John 4:7

Principle: The strength of community comes from relationships.

When a polymer cross-links, the physical properties can change dramatically. Without the cross-links, some polymers are very fluid. But once the linkages are in place, this same substance can be a rigid solid. Epoxy is like this. The strength of the cross-linked polymer comes from both the number of linkages and the strength of the linkages. In Peter’s illustration of the house given above, the building materials are added to one another to make a structure. If the structure were of stone, the stones would be cut and fit into place so that they interlocked with each other. Once interlocked, the stone walls could stand for centuries, or even longer.

Communities gain their strength from relationships. In some communities, people know each other for a lifetime. They have been through storms and through stress. They may have helped and served each other. All this makes a relational linkage between the people. These are bonds of community. The same can be true in the family as well. In the church,

there is an additional element that binds people together. The Holy Spirit acts as an agent to bring the community together in love. Again, it is through common ministry together or by serving one another that bonds of friendship and love are established. Sometimes these bonds are so strong that they can be sensed by those who enter from the outside. Those who don’t have love in their lives may be drawn to this love. Sharing why we love is a wonderful way to bring people into the kingdom of God.

Procedure

Note: Sodium tetraborate is slightly toxic (it is used as a laundry detergent enhancer). Do not ingest the slime, and wash your hands after handling it.

Note: Heat the water before you add the PVA.

Note: You may use the digital balance to measure the reagents.

1. Put 50 mL of hot water in the 150-mL beaker (or heat the water to about 85° C in a microwave). Use the 1-mL measuring spoon to add 3 scoops (about 2.1 g) of polyvinyl alcohol (*PVA*) to the 50 mL of hot water in the beaker. Stir this with the stir stick to dissolve the *PVA*. Allow this to cool until it is comfortable to handle.

Note: It might take some time to dissolve all the PVA, and you may need to reheat the water.

2. Observe and record the properties of the *PVA* solution (question 1 in the questions section). Add a drop or two of food coloring if you desire to have the slime colored.

3. While the *PVA* solution is cooling, stir one half ($\frac{1}{2}$) to three fourths ($\frac{3}{4}$) of a scoop (about 0.4 - 0.6 g) of sodium tetraborate into 10 mL of warm water in the 30-mL beaker.

4. Stir the sodium tetraborate solution

into the *PVA* solution. While stirring, lift the stir stick out of the solution several times to observe the viscosity and other property changes. Record this in the questions section.

5. It may take several minutes for the cross-linking to occur and for the material to become more solid. This result is slime.

6. Put the slime on a paper plate or other clean surface.

7. Remember to clean up.

6. A non-Newtonian fluid is a fluid in which the viscosity changes with the applied strain rate. If pressure is applied quickly, it acts differently than if pressure is applied slowly.

a.) Pull the material apart quickly. What happens? Put the material back together and slowly pull it apart. What is the difference? What effect do the molecular mass of the *PVA* and hydrogen bond cross-links have on its physical properties?

b.) Roll a small piece of slime into a ball and drop it. What happens? Place that ball onto the counter for a few minutes. What happens? Is slime a non-Newtonian fluid? Why or why not? You might want to do more research about non-Newtonian fluids on the Internet.

7. The cross-linked polymer initially contains a high percentage of water (about 96% water). How do the properties of the material change as the water evaporates out of it? You may want to leave a piece of the material uncovered overnight to observe how it changes when the water it contains evaporates.

